



Kongeriget Danmark

Patent application No.: PA 1999 01395

Date of filing: 30 September 1999

Applicant: Rolf Jess Jørgensen
Rytterhaven 48
DK-3520 Farum

This is to certify the correctness of the following information:

The attached photocopy is a true copy of the following document:

- The specification, claims, abstract and drawings as filed with the application on the filing date indicated above.



**Patent- og
Varemærkestyrelsen**
Erhvervsministeriet
TAASTRUP 26 April 2000

Lizzi Vester

Lizzi Vester
Head of Section

A METHOD OF PREVENTING HYPOCALCEMIA IN LACTATING ANIMALS AND COMPOSITIONS USED THEREIN

5 FIELD OF THE INVENTION

The present invention provides a method of preventing hypocalcemia in lactating animals and in particular there is provided compositions which can be administered to the animals during the dry period to prevent hypocalcemia and milk fever.

10

TECHNICAL BACKGROUND AND PRIOR ART

Hypocalcemia is a metabolic disease of lactating animals, such as dairy cows, and occurs as subclinical hypocalcemia around calving and in early lactation and in severe cases
15 lactating animals contract clinical milk fever (parturient paresis). The lactation cycle of a dairy cow extends for approximately 10 month of the year immediately subsequent to calving. Lactation by dairy cows is at a minimum during the two month immediately prior to calving, and the dairy cows are normally not milked during this "dry period". The act of calving induces lactation in the dairy cow, and milk production is at a peak by 4-6 weeks
20 thereafter.

During the dry period, the dairy cow has a low demand for calcium, and is thus able to absorb sufficient calcium via the digestive tract by simple passive diffusion. During lactation the demand for calcium is high, which leaves the cow in a negative calcium
25 balance, which results in the activation of the vitamin D-dependent complex mechanisms. The lactating cow is thus capable of increasing the utilisation of the feed ration calcium by increasing absorption from the digestive tract and by resorbing calcium from the skeletal system via the blood. By these active mechanisms the lactating cow is capable of covering up to 80% of its demand for calcium during lactation (Klooster, 1976). Therefore,
30 lactating animals are capable of preventing hypocalcemia to occur due to these powerful natural defence mechanisms almost regardless of the calcium intake via the feed.

However, during the dry period these natural defence mechanisms are deactivated, as the demand for calcium is low. Upon calving and onset of lactation there is an immediate
35 demand for high levels of calcium to be supplied via the blood circulation to the mammary

glands for milk production. The calcium regulating mechanisms are not prepared for this sudden demand for calcium in the high yielding dairy cow, and it must thus contract hypocalcemia and milk fever. Symptoms of milk fever, such as lowering of body temperature, usually become manifest within one day before and four days after calving.

5

Extensive research has been conducted to develop methods for preventing or treating milk fever. It has been observed that the risk for milk fever is reduced when the acid-base balance of the dairy cow is manipulated by acidifying dietary ration composition (Dishington, 1982). Presently, a method based on this principle is widely used in the USA.

10 However, acidification of dairy cows is unnatural, since urine from ruminants normally is alkaline. Furthermore, overdosage has been reported.

Another approach of preventing milk fever is the peroral drenching with calcium chloride around calving (Jørgensen, 1990). The effect of this method is interpreted as a simple
15 calcium diffusion through the wall of the forestomach and through the wall of the small intestine. However, there are several drawbacks associated with this method. The dairy cows have to be handled and dosed individually, which involves the above mentioned problems. Furthermore, the precise calving time of the cow must be known, because the first drenching is recommended to be carried out a few hours before calving.

20

Injection of a synthetic D-vitamin preparation in large doses has been suggested for the prevention and/or treatment of milk fever. Although this method is documented to be effective, it is not allowed in many countries. Another suggestion involves continuous milking, which will prevent milk fever, as the calcium regulating mechanisms are
25 constantly activated. However, this method is only applicable to single cows which are known or suspected to contract milk fever, but it is not feasible at a herd level. Furthermore, it has been observed that the risk of milk fever can be reduced by feeding the dairy cows a dietary ration having a high calcium content prior to calving.

30 Ramberg et al. (1970) observed that milk fever can be controlled and/or eliminated by administering a low calcium dietary ration to dairy cows during the weeks immediately prior to calving. By feeding a low calcium dietary ration, the cow is left in a negative calcium balance, which results in the activation of its natural calcium regulating mechanisms. Unfortunately, there are many problems associated with this method. For
35 example, most of the cattle feed, such as alfalfa and other legumes, contain too high

1 v ls of calcium to bring the dry cow in a negative calcium balance. Furthermore, the method pres nts a substantial problem when dealing with a large herd. To solve th above first mentioned problem, US 3,908,020 discloses a complete, calcium-limited feed ration. However, this method implies the purchase of rather expensive feed, which is not
5 feasible under commercial dairy farming conditions, especially when the farmer is using his own grown crops for feeding.

In view of the foregoing it is evident that these currently used methods of preventing hypocalcemia involve several problems in modern farming. It is thus an important
10 objective of the present invention to provide a novel composition and method to prevent hypocalcemia in lactating animals. The disclosed method has the advantage that its effect is based on the powerful, natural calcium regulating mechanisms of animals and at the same time it is applicable under commercial dairy farming conditions.

15

SUMMARY OF THE INVENTION

Accordingly, it is the primary objectives of the present invention to provide a novel method of preventing hypocalcemia in lactating animals and a composition which is useful in such
20 a method.

Thus, in a first aspect, the invention pertains to a method of preventing hypocalcemia in a lactating animal, comprising administering to the animal during at least part of the dry period an effective amount of a compound which reduces absorption of calcium from
25 drinking water and/or from the ration of said animal.

There is also provided a composition for preventing hypocalcemia in a lactating animal, comprising an encapsulated compound which reduces the absorption of calcium from the drinking water and/or from the ration of said animal.

30

In a further aspect, the present invention relates to the use of the above composition for the prevention of hypocalcemia in a lactating animal, comprising administering to the animal during at least part of the dry period an effective amount of a compound which reduces the absorption of calcium from the drinking water and/or from the ration of the
35 animal.

In yet another aspect the invention pertains to a method of using the above composition for the prevention of hypocalcemia in a lactating animal, the method comprising administering to the animal during at least part of the dry period an effective amount of a
5 compound which reduces the absorption of calcium from the drinking water and/or from the ration of the animal according to the above method.

In a still further aspect, the invention relates to the use of a calcium-binding compound or a composition in the manufacturing of a medicament for preventing hypocalcemia in a
10 lactating animal.

DETAILED DISCLOSURE OF THE INVENTION

15 It is an essential feature of the method for preventing hypocalcemia in lactating animals which is provided herein that the biological effect of the compound used in the method is capable of stimulating or triggering the natural calcium regulating defence mechanism of the animal. Through this stimulating ability on the calcium metabolism before the onset of lactation, the used compound will render the animal well prepared to resist any
20 threatening parturient hypocalcemia.

Thus, the method according to the invention comprises administering to the animal during at least part of the dry period an effective amount of a compound which reduces absorption of calcium from drinking water and/or from the ration of said animal. As used
25 herein, the expression "lactating animal" designates the mammalian class of the animal kingdom such as e.g. cow, pig, horse, cat, sheep, goat, buffalo, camels, monkey or man. Thus, the term "ration" relates both to "food" and "feed" given prior to the onset of lactation of said animal.

30 An advantageous feature of the method according to the invention is that the precise parturition time of the animal does not need to be known, as the compound can be given to the animal at any time during the dry period, such as at least about 1 to 4 weeks prior expected parturition, including at least about 2 to 3 weeks prior to expected parturition, and in amounts of about 10 to 1000 grams per animal per day depending on the body
35 weight of the animal.

In presently preferred embodiments, the method according to the invention comprises administering to the animal during at least part of the dry period an effective amount of a calcium-binding compound. It will be understood that the expression "calcium-binding
 5 compound" refers to compounds which are capable of binding free (ionised) calcium in aqueous solution.

Calcium-binding compounds which are useful in the above method can be any compounds which are capable of binding free calcium in the animal gastro-intestinal tract
 10 and thus trigger the natural calcium regulating defence mechanism of the animal. Such compounds include compounds that are either digestible or non-digestible in the rumen selected from the group consisting of oxalic acid, sodium oxalate, phytic acid, a phytate, a clay mineral including zeolite, sodium diethylene acetic acid, ethylenediaminetetraacetic acid (EDTA) and its sodium salts Na_2EDTA and Na_4EDTA , trisodium nitrilotriacetate
 15 monohydrate, trisodium nitriloacetate, pentasodium diethylenetriaminepentaacetate, trisodium N-hydroxyethylethylenediaminetriacetate, citric acid, a citrate, a polyphosphate, a tripolyphosphate, an orthophosphate and a cellulose phosphate and calcium-free derivatives of any such compounds.

20 In a presently preferred embodiment, the method according to the invention comprises administering to the animal during at least part of the dry period an effective amount of a compound which has a competitive effect on the absorption of calcium from the drinking water and/or from the ration of said animal through the wall of the gastro-intestinal tract. It is contemplated that such compounds will compete with the calcium of the dry animal
 25 dietary ration by being preferred by the calcium absorbing mechanism. Examples of such compounds are zinc compounds including zinc oxide, ZnCl_2 and ZnSO_4 .

It is, as mentioned above, of considerable commercial interest to provide a method of preventing hypocalcemia in lactating animals which is applicable under commercial
 30 farming conditions. In a highly convenient embodiment of the method according to the invention, the compound used in the method is contained in a composition which is in the form of a premix, a liquid or a powder. It is contemplated that such a composition can easily be administered to the dry animal perorally in a specific embodiment, be added to the drinking water and/or the ration of the dry animal.

35

In a further aspect of the invention there is provided a composition, which is used in the above method for preventing hypocalcemia in a lactating animal. The composition comprises, in encapsulated form, a compound which reduces the absorption of calcium from the drinking water and/or from the ration of said animal.

5

In an advantageous embodiment of the composition according to the invention the encapsulated compound is any of the above calcium-binding compounds. It will be understood that it is possible to have a mixture of two or more calcium-binding compounds in the same composition.

10

In one useful embodiment of the composition according to the invention, the compound has a competitive effect on the absorption of calcium from drinking water and/or from the ration of the dry animal. Typical examples of such compounds are zinc compounds such as e.g. zinc oxide, ZnCl_2 or ZnSO_4 .

15

As mentioned above, the compounds which are used in the above compositions are encapsulated by any appropriate encapsulating material. In specific embodiments of the invention, a useful compound for the encapsulation is a compound selected from the group consisting of a fat, a non-calcium derivative of a fat such as a soap and a stearate, a protein, a polysaccharide, a cellulose and a derivative of any such compound, a gum, a glycol and gelatine.

20

In an interesting embodiment, the composition according to the invention comprises compounds which are encapsulated by a calcium-free membrane material, which at the body temperature of the lactating animal is solid at a pH value above 4.0 but which under these conditions dissolves at pH below 4.0. It is contemplated that compounds encapsulated in such a membrane material can be transported through the rumen of a ruminant without being dissolved, and thus is not dissolved until it arrives in the gastrointestinal system after passage of the forestomachs of the ruminant.

30

In a highly convenient embodiment of the invention, the composition is in the form of a premix, a liquid and a powder, or it in the form of a dragee, a tablet, a pill, or a capsule. A composition provided in such a form will be advantageous from a users point of view, since it can easily be administered to the dry animal. In a specific embodiment, the composition is added to or mixed into the drinking water and/or in the ration of the

35

lactating animal. It is evident that such a composition is very useful, when farmers use their own grown crops as feed.

In useful embodiments, the composition according to the invention is a composition
 5 comprising the above compounds and at least one further ingredient such as a vitamin, a mineral and a carrier. A useful example of such a carrier is e.g. bran. Further ingredients can be selected from a sugar, a stabilising agent and a colouring agent. Additionally, the composition is calcium-free and non-acidifying, i.e. the composition is not causing biological acidification of the animal's urine.

10

In further aspects, the invention relates to the use of a composition according to the invention, comprising administering to the animal during at least part of the dry period an effective amount of a compound which reduces the absorption of calcium from the drinking water and/or from the ration of the animal.

15

In a still further aspect the invention relates to the use of a composition according to the present invention for the prevention of hypocalcemia in a lactating animal.

Furthermore, there is provided a method of using a composition according to the invention
 20 for the prevention of hypocalcemia in a lactating animal, the method comprising administering to the animal during at least part of the dry period an effective amount of a compound which reduces the absorption of calcium from the drinking water and/or from the ration of the animal according to the method of the present invention.

25 In addition, the present invention relates to the use of a calcium-binding compound or a composition according to the invention in the manufacturing of a medicament for preventing hypocalcemia in a lactating animal.

The invention will now be described in further details in the following non-limiting
 30 examples and the drawings wherein

Fig. 1 shows the response of three cows (Nos. 169, 170 and 168) to repeated short-term periods of hypocalcemia induced by EDTA test infusions. The other three cows (Nos. 172, 175 and 176) were dosed daily with anionic salts prior to the EDTA-testing. Cal = ionised
 35 calcium in the blood;

Fig. 2 shows the blood calcium response to oral administration of zinc oxide in pairs of lactating as well as non-lactating cows. A total of four treatments were given at one-week intervals. Arrows indicate time of treatment;

5

Fig. 3 shows the blood calcium concentration of the lactating cows given four doses of zinc oxide at one-week intervals, as well as of the non-lactating, untreated cows. Arrows indicate time of treatment;

10 Fig. 4 shows the blood calcium response of two lactating cows to the four doses of zinc oxide expressed in terms of the extent of the drop in blood calcium;

Fig. 5 shows the blood calcium of five dairy cows in late lactation. The graph shows the response, in form of suppression of the daily increase of blood calcium, during

15 administration of 2x500g of a synthetic zeolite A (sodium aluminium silicate) per day, and a late response in form of "overshooting" at 24 hours after the last administration and onwards; and

Fig. 6 shows the blood calcium of four non-pregnant dry cows: Two untreated controls

20 and two treated with 2 kg zeolite A (sodium aluminium silicate) once a day on day 8, 9, 10 and 11.

EXAMPLE 1

25

Repeated EDTA-infusions in non-lactating (dry) cows

1.1. Introduction

30 It is known that hypocalcemia can be induced experimentally by slow intravenous infusion of EDTA. Such induced hypocalcemia is performed for a variety of reasons, as reviewed by Jørgensen et al. (1999).

35

1.2. Materials and methods

To investigate the induction of hypocalcemia by intravenous infusion of EDTA in cows, 6 non-lactating (dry) cows were used in this study. Three of these cows were untreated (cow Nos. 168, 169 and 170) and the other three cows (cow Nos. 172, 175 and 176) had been subjected to a pre-treatment consisting of daily doses of anionic salts (Wang and Beede, 1992). This pre-treatment is known to be effective in preventing milk fever. The latter cows were rumen fistulated to facilitate precise daily dosages of the anionic salts.

All cows were tested for resistance to hypocalcemia by the EDTA-infusion technique. A total of 8 EDTA-tests were performed per cow according to the standardised method of Mellau et al. (1998), at 10 days intervals. With the standardised test, infusion of EDTA was performed until the cow went down involuntarily, due to muscle weakness caused by the induced hypocalcemia. The time taken from this point and until the cow regained normal blood calcium concentration, which is defined as an ionised blood calcium value above 1 mmol/l, was recorded. This time is considered to represent the cows' calcium mobilisation rate and therefore also the cows' ability to resist hypocalcemia. The infusion periods varied between individual cows, but were typically in the range of 2-4 hours. The total time exposed to hypocalcemia during one test, i.e. during infusion plus during recovery, was in the range of 4-8 hours.

1.3 Results

It is obvious from Fig. 1 that the time it took to regain normal blood calcium concentration greatly improved in the three untreated control cows after three infusions in terms of reduced recovery time and therefore a fall in the graphs for individual cows indicated increased resistance to hypocalcemia. For comparison, the results of three other cows tested in the same way in the same experiment, but after a 10-day period of daily anion treatment prior to EDTA-testing, are also shown. As would be expected the recovery time for these cows was relatively short already at the first EDTA-test due to the treatment prior to the test period.

1.4 Discussion

Hypocalcemic episodes of a few hours duration, induced three times at 10-day intervals, resulted in shortened recovery periods at later similar challenges. The effect was of the same magnitude as that recorded after daily anion supplementation, which is a well-known milk fever preventive method.

5

EXAMPLE 2

Repeated zinc oxide administration in cows

10

2.1 Introduction

Until now, the only known way of inducing hypocalcemia via the oral route (by mouth) has been that recorded as an undesired side-effect of administration of zinc in the form of zinc oxide to cattle and small ruminants (Smith et al. 1984). Such treatment is given against facial eczema in sheep and cattle. The possible invasive effect of repeated challenges towards resistance to hypocalcemia was examined in this study involving lactating as well as non-lactating (dry) cows.

20 2.2 Materials and methods

Six dairy cows were used in this study: Two lactating cows and two dry cows were treated with zinc oxide by mouth at 120mg/kg body weight, whereas two lactating cows served as untreated controls. Four doses of zinc oxide were given to the treated cows at one-week intervals. Venous blood samples were taken and serum stored at -37°C for later total calcium determinations. For this purpose blood samples were taken daily from all six cows one hour after morning feeding. Extra samples were taken 5, 11, 23 and 35 hours after zinc oxide administration.

30 2.3 Results

Fig. 2 shows the response in serum calcium concentration of the lactating cows as well as of the non-lactating cows treated with the same amount of zinc oxide per kg of body weight. It is obvious that the most pronounced response was obtained in the lactating cows, followed by the dry cows. In the lactating cows the resulting hypocalcemia lasted

about 24 hours with a minimum of 1.4 - 1.6 mmol/l occurring approx. 24 hours after administration. The minimum values were lower than the lowest accepted value for normal serum calcium concentration of 2.17 mmol/l (Rosol et al. 1995). In Fig. 3 the response of the lactating cows is shown more clearly on the background of the blood values of the untreated non-lactating controls. Fig. 4 shows that the response of the lactating cows to the zinc oxide doses changed with time, i.e. with the number of doses in that the cow's calcium balance became less and less affected.

2.4 Discussion

10

This experiment shows that hypocalcemic episodes of approximately 24 hours duration could be produced repeatedly in dairy cows by oral zinc oxide administration at one-week intervals. Provided it is accepted that the graphs in Fig. 4 has a downwards course (in statistical terms: that they are different from horizontal) it shows that the cow's resistance to the induced hypocalcemia challenge increased with the number of zinc oxide induced hypocalcemic periods.

EXAMPLE 3

20

Repeated administration of synthetic sodium aluminium zeolite A to lactating cows

3.1 Introduction

So-called calcium binders are widely used in various industries such as the soap and detergent industry, or as water softeners in water treatment plants. Many natural minerals of the zeolite family as well as synthetic zeolites are strong cation exchangers and are used as such for the attraction of calcium in the form of Ca^{2+} in exchange of other cations like sodium (Na^{2+}). In the following experiment a group of dairy cows were given such a zeolite and their serum calcium was monitored. The purpose was to test any direct effect on serum calcium during the drenching period as well as any delayed effect likely to be linked to resistance to hypocalcemia. In this experiment lactating dairy cows were used as a model for non-lactating (dry) cows.

3.2 Materials and methods

35

Five Red Danish Dairy cows in late lactation were used. The cows yielded approximately 20 kg of milk per day and were milked only once daily, i.e. in the morning. After an initial baseline period each cow received 0.5 kg of zeolite by stomach tube twice a day for two consecutive days. On the third day cows were only treated in the morning. In this way
 5 each cow received a total dosage of 2.5 kg. The zeolite used was a synthetic sodium aluminium silicate (Dedima 100; Degussa-Hüls AG, Germany). All cows were bled twice a day: at 10 a.m. and at 22 p.m. Serum was stored at -18°C and later analysed for total calcium concentration.

10 3.3 Results

Fig. 5 shows the average serum calcium response of the five cows. It can be seen that the serum calcium baseline values before administration fluctuated daily with comparatively low values in the morning followed by an increase during the day. During
 15 zeolite administration there was a repression in the increase in serum calcium during the day, as compared to before administration. From 24 hours after last administration and onwards full daily fluctuations were restored, but they now took place at a higher level as compared to the initial baseline level.

20 3.4 Discussion

This is a first testing of Dedima 100 in cows and it shows that at this dose level blood calcium is repressed during administration and that the animals respond to this by elevated calcium level after the administration period. The latter phenomenon is known
 25 from other descriptions of hypocalcemia and has been termed "overshooting" by Hove (1987) who describes it as a response to released parathyroid hormone (PTH) which is part of the organism's immediate defence mechanism against hypocalcemia.

30 EXAMPLE 4

Repeated administration of synthetic sodium aluminium zeolite A to dry cows

4.1 Introduction

35

The present experiment was performed with dry, non-pregnant cows as a model for dry, pregnant cows. Non-pregnant dry cows are expected to be less sensitive to hypocalcemia than pregnant dry cows because the former has the lowest requirement for calcium.

5 4.2 Materials and methods

Four non-pregnant dry dairy breed cows were used. Two of these were treated with Dedima 100 at a single dose of 2 kg/day for 4 days, whereas the other two served as untreated controls. Venous blood samples were taken daily from all four cows at 10.00 hours, i.e. two hours after morning feeding. Extra blood samples were taken on the days of administration at 22.00 hours.

4.3 Results

15 When compared with the untreated controls, a decrease in the concentration of the blood calcium was seen in the two experimental cows following the experimental treatment. As it can be seen from Fig. 6, in one cow (the zeolite 1 - cow) this decrease in the concentration of the blood calcium was only slight and lasted only 1 or 2 days. In the other cow (the zeolite 2 - cow) a more obvious decrease in blood calcium concentration was
20 recorded. It was pronounced 2 days after the first dose was given and had reached 1.8 mmol/l by the third day after first administration and remained subnormal until 3 days after the last dose was given. There was a reduction in appetite in both the experimental cows and transient constipation was recorded in the zeolite 2 - cow.

25 4.3 Discussion

This experiment demonstrates that hypocalcemia can be induced in a non-pregnant dry cow by the oral administration of a zeolite for 4 days. It also shows, that individual variation among such cows can be markedly. The recorded difference was similar to the
30 difference seen in the same cows in a previous study (Example 2) with zinc oxide.

REFERENCES

- Dishington, I.W. and Bjoernstad, J. 1982. Prevention of milk fever by dietary means. The effect of a concentrate fortified with mineral salts. *Acta vet. scand.*, **23**, 336-343.
- 5
- Hove, K. 1987. Milk fever prevention and calcium homeostasis around calving in the Dairy cow. *Agripractice; Irish Veterinary news*, March 1987:24-27.
- Jørgensen, R.J., Basse, A. and Aslan, V. 1990. Sequelae to oral calcium chloride gel dosing of cows. *Proceedings, XVI World Buiatrics Congress*, 13.-17. August 1990, Salvador, Bahia, Brazil, pp. 511-515.
- 10
- Jørgensen, R. J., Nyengaard, N. R., Daniel, R. C. W., Mellau, L. S. B., Enemark, J. M. D. 1999. Induced hypocalcaemia by Na₂EDTA infusion. A review. *Journal of Veterinary medicine A*, Vol. **46**(7):389-452.
- 15
- Mellau, L. S. B., Jørgensen, R. J., Enemark, J. M. D. 1998. Standardization and interpretation of experimental (Na₂EDTA induced) hypocalcaemia in cows. 10th ICPD (International Conference on Production Diseases in Farm Animals), Utrecht, p. 238.
- 20
- Ramberg, C.F., Phang, J.M. and Kronfeld, D.S. 1970. A compartmental model of calcium metabolism in cows. In: *Parturient hypocalcemia*. Academic Press, New York, pp. 119-134.
- 25
- Rosol, T. J., Chew, J. D., Nagode, L. A., Capen, C. C. 1994. Pathophysiology of calcium metabolism. *Veterinary Clinical Pathology*, **24**:49-63.
- Smith, B. L., Collier, A. J., Lawrence, R. J., Towers, N. R. 1984. Hypocalcaemia associated with high dose rates of zinc oxide to lactating dairy cows. *New Zealand Veterinary Journal*, **32**:48-50.
- 30
- Van't Klooster, A.Th. 1976. Adaptation of calcium absorption from the small intestine of dairy cows to changes in the dietary calcium intake and at the onset of lactation. *Z. Tierphysiol., Tierernährg. u. Futtermittelkde*, **37**:169-182.
- 35

Wang, C. and Beede, D. K. 1992. Effect of ammonium chloride and sulphate on acid-base status and calcium metabolism of dry cows. *Journal of Dairy Science*, **75**:820-828.

CLAIMS

1. A method of preventing hypocalcemia in a lactating animal, comprising administering to the animal during at least part of the dry period an effective amount of a compound which
5 reduces absorption of calcium from the drinking water and/or from the ration of said animal.
2. A method according to claim 1, comprising administering to the animal during at least part of the dry period an effective amount of a calcium-binding compound.
- 10 3. A method according to claim 2 wherein the calcium-binding compound is selected from the group consisting of oxalic acid, sodium oxalate, phytic acid, a phytate, a clay mineral including zeolite, ethylenediaminetetraacetic acid (EDTA) and its sodium salts Na₂EDTA and Na₄EDTA, trisodium nitrilotriacetate monohydrate, trisodium nitriloacetate,
15 pentasodium diethylenetriaminepentaacetate, trisodium N-hydroxyethylethylenediaminetriacetate, citric acid, a citrate, a polyphosphate, a tripolyphosphate, an orthophosphate and a cellulose phosphate and a calcium-free derivative of any such compounds.
- 20 4. A method according to claim 1, comprising administering to the animal during at least part of the dry period an effective amount of a compound which has a competitive effect on the absorption of calcium from the drinking water and/or from the ration of said animal.
5. A method according to claim 4 wherein the compound is a zinc compound including
25 zinc oxide, ZnCl₂ and ZnSO₄.
6. A method according to claim 1 wherein the compound is contained in a composition which is in the form of a premix, a liquid and a powder.
- 30 7. A method according to claim 6 wherein the composition is added to the drinking water and/or to the ration of the dry animal.
8. A composition for preventing hypocalcemia in a lactating animal, comprising, in encapsulated form, a compound which reduces the absorption of calcium from the
35 drinking water and/or from the ration of said animal.

9. A composition according to claim 8 where the compound is a calcium-binding compound.
- 5 10. A composition according to claim 9 where the calcium-binding compound is selected from the group consisting of oxalic acid, sodium oxalate, phytic acid, a phytate, a clay mineral including zeolite, ethylenediaminetetraacetic acid (EDTA) and its sodium salts Na_2EDTA and Na_4EDTA , trisodium nitrilotriacetate monohydrate, trisodium nitriloacetate, pentasodium diethylenetriaminepentaacetate, trisodium N-hydroxyethyl-
- 10 ethylenediaminetriacetate, citric acid, a citrate, a polyphosphate, a triphosphate, an orthophosphate and a cellulose phosphate and a calcium-free derivative of any such compounds.
11. A composition according to claim 8 where the compound has a competitive effect on
- 15 the absorption of calcium from drinking water and/or from the ration of the animal.
12. A composition according to claim 11 where the compound is a zinc compound including zinc oxide, ZnCl_2 and ZnSO_4 .
- 20 13. A composition according to claim 8 where the compound is encapsulated by a compound selected from the group consisting of a fat, a non-calcium derivative of a fat such as a soap and a stearate, a protein, a polysaccharide, a cellulose, a gum, a glycol, gelatine and a derivative of any such compound.
- 25 14. A composition according to claim 8 where the compound is encapsulated by a calcium-free membrane material, which at the body temperature of the lactating animal is solid at a pH value above 4.0 but which dissolves at pH below 4.0.
15. A composition according to claim 8 where the composition is in the form of a premix, a
- 30 liquid and a powder.
16. A composition according to claim 15 where the composition is added to the drinking water and/or to the ration of the dry animal.

17. A composition according to claim 8 where the composition comprises at least one further ingredient such as a vitamin, a mineral or a carrier.

18. A composition according to claim 8 where the composition is calcium-free and non-
5 acidifying.

19. Use of a composition according to any of claims 8-18 for the prevention of hypocalcemia in a lactating animal, comprising administering to the animal during at least part of the dry period an effective amount of a compound which reduces the absorption of
10 calcium from the drinking water and/or from the ration of the animal.

20. A method of using a composition according to any of claims 8-18 for the prevention of hypocalcemia in a lactating animal, the method comprising administering to the animal during at least part of the dry period an effective amount of a compound which reduces
15 the absorption of calcium from the drinking water and/or from the ration of the animal.

21. Use of a calcium-binding compound or composition according to any of claims 8-18 in the manufacturing of a medicament for preventing hypocalcemia in a lactating animal.

ABSTRACT

A METHOD OF PREVENTING HYPOCALCEMIA IN LACTATING ANIMALS AND
COMPOSITIONS USED THEREIN

5

Method of preventing hypocalcemia and milk fever (parturient paresis) in lactating animals comprising administering to the animal during the dry period a compound which reduces absorption of calcium from the drinking water and/or from the ration of said animal. The method is based on the powerful, natural calcium regulating mechanisms and is
10 applicable under commercial farming conditions.

1/6

Modtaget PD
30 SEP. 1999




Fig. 1

2/6




Fig. 2

3/6




Fig. 3

4/6




Fig. 4

5/6




Fig. 5

6/6




Fig. 6